



University of Groningen

Cohomology and the Resolution of the Nilpotent Variety

Hesselink, Wim H.

Published in:
Mathematische annalen

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1976

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Hesselink, W. H. (1976). Cohomology and the Resolution of the Nilpotent Variety. *Mathematische annalen*, 223(3), 249-252.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Cohomology and the Resolution of the Nilpotent Variety

Wim H. Hesselink

Department of Mathematics, University of Groningen, Postbox 800, Groningen, The Netherlands

1. Let G be a split reductive linear algebraic group over a field k of characteristic zero. Consider the variety N of the nilpotent elements of the Lie algebra \mathfrak{g} of G . It is a normal variety cf. [14] Theorem 16. It is isomorphic to the variety of the unipotent elements of G , cf. [17]. The theorem of Brieskorn-Steinberg-Tits states that the rational singularities are dense in the singular locus of N , see [1] and [18] (3.10). Here we shall prove that N has only rational singularities, cf. [12] p. 50, i.e. we prove

Theorem A. *There exists a proper birational morphism $\tau: Y \rightarrow N$ such that Y is smooth over k , that $\tau_*(\mathcal{O}_Y) = \mathcal{O}_N$ and $R^p\tau_*(\mathcal{O}_Y) = 0$ for $p \geq 1$.*

This theorem admits a generalization which will be stated and proved in Section 5. In the complex analytic situation the same assertions follow by Theorem 5 of [3] exp. II. For rational singularities in that case see [2]. In [11] we investigated the local structure of N for the classical groups.

2. By [17] (2.2) we may assume that G is semi-simple and simply connected. Let G be split with respect to a maximal torus T and a Borel group B .

If E is a finite dimensional vector space over k and $\varrho: B \rightarrow GL(E)$ is a morphism of algebraic groups then $E = (E, \varrho)$ is called a B -module. The contracted product $G \times^B E$ is defined as the quotient of $G \times E$ under the right action of B given by $(g, e)b = (gb, \varrho(b^{-1})e)$ for all $g \in G, e \in E, b \in B$. The morphism $\psi: G \times^B E \rightarrow G/B$ given by $\psi(g, e)B = gB$, is a vector bundle over G/B . The sheaf of sections of ψ is a locally free $\mathcal{O}_{G/B}$ -module. It is denoted by $\mathcal{L}(E)$. See [7] p. 55, 56.

Let \mathfrak{u} be the Lie algebra of the unipotent part U of B . As \mathfrak{u} is a B -module we can define $Y = G \times^B \mathfrak{u}$. The adjoint action $\text{Ad}: G \times \mathfrak{g} \rightarrow \mathfrak{g}$ induces a surjective morphism $\tau: Y \rightarrow N$. Since $\psi: Y \rightarrow G/B$ is a vector bundle, Y is an irreducible smooth variety. The morphism τ is easily identified with the G -equivariant proper morphism τ considered in [17] (the proof of 2.1), which is birational, cf. [17] (2.4). Since N is normal, it follows that $\mathcal{O}_N = \tau_*(\mathcal{O}_Y)$. By [10] (1.4.11) it suffices now to prove that $H^p(Y, \mathcal{O}_Y) = 0$ for all $p \geq 1$. This is a special case of the corollary in Section 5.

Remark. The action of B on u is not completely reducible, so we cannot apply the theorem in [13].

3. Let $X(T)$ be the character group of T . Let R be the root system of G with respect to T and let W be the Weyl group. Conform [5] and [6] a root α is called positive if its eigenspace \mathfrak{g}_α is not contained in u . The set of positive roots is denoted by R_+ . For each root α we have an associated co-root α^* in the dual \mathbb{Z} -module $X(T)^*$. A weight $\chi \in X(T)$ is called regular (resp. dominant) if we have $\langle \alpha^*, \chi \rangle \neq 0$ (resp. $\langle \alpha^*, \chi \rangle \geq 0$) for all $\alpha \in R_+$. The set of dominant weights is denoted by $X(T)_+$. If V is a subset of R we write $|V| = \sum_{\alpha \in V} \alpha$. The number of elements of V is denoted by $\#(V)$. The weight ϱ is defined by $2\varrho = |R_+|$. It is well known that $\varrho + X(T)_+$ is the set of regular dominant weights. The length of $w \in W$ is denoted by $n(w)$, cf. [6].

If E is a B -module we write $H^p(E) = H^p(G/B, \mathcal{L}(E))$. As \mathcal{L} is an exact functor, the functors H^p form an exact delta-functor from the category of B -modules to the category of G -modules. For $\chi \in X(T)$ let $E(\chi)$ be the one-dimensional B -module corresponding to the induced morphism $B \rightarrow T \rightarrow Gl(1)$. We shall use the following version of Bott's theorem, cf. [6].

If $H^p(E(\chi)) \neq 0$, then $\chi + \varrho$ is regular and $p = n(w)$, where w is the unique element of W such that $w(\chi + \varrho)$ is dominant (and regular).

4. Definition. If $\mu \in X(T)$, let $p(\mu)$ be the maximal value of $n(w) - \#(V)$, where $w \in W$ and V is a subset of R_+ such that $w(\mu + \varrho - |V|)$ is dominant and regular.

Lemma. Let $\mu \in X(T)$.

(a) $p(\mu)$ is the maximal value of $\#(V \cap -wR_+) - \#(V \cap wR_+)$, where $w \in W$ and V is a subset of R_+ such that $w(\mu) - |V|$ is dominant.

(b) We have $0 \leq p(\mu) \leq \#(R_+)$.

(c) If $\mu = 0$ then $p(\mu) = 0$.

Proof. Consider subsets P of R satisfying $P \cap -P = \emptyset$ and $P \cup -P = R$. The relations $V = R_+ \cap P$, $P = V \cup -(R_+ \setminus V)$ define a one-to-one correspondence between these subsets of R and the subsets V of R_+ . The natural action of W on the collection of the subsets P induces an action of W on the power set of R_+ , which is given by

$$w * V = R_+ \cap (wV \cup -w(R_+ \setminus V)).$$

If V corresponds to P then $\varrho - |V| = -\frac{1}{2}|P|$. This implies

$$w(\varrho - |V|) = \varrho - |w * V|.$$

It follows that $w(\mu + \varrho - |V|)$ is dominant and regular if and only if $w(\mu) - |w * V|$ is dominant. As $n(w) = \#(R_+ \cap -wR_+)$ we have

$$n(w) - \#(V) = \#((w * V) \cap -wR_+) - \#((w * V) \cap wR_+).$$

Now (a) follows immediately. (b) is a consequence of (a). If V is a non-empty subset of R_+ then $-|V|$ is not dominant. So (c) follows from (a). Compare [4] and [15] (2.13).

Remark. It seems that $\mu \in \varrho + X(T)_+$ implies $p(\mu) = 0$. If R is a root system of type A_l and α is the highest root, then we have $\alpha \in X(T)_+$ and $p(\alpha) \geq l - 4$.

5. The notations are as before. G is semi-simple and simply connected. Let \mathfrak{p} be a linear subspace of \mathfrak{g} which is B -invariant and contains u . Consider $Z = G \times^B \mathfrak{p}$ and the canonical morphism $\psi: Z \rightarrow G/B$.

Theorem B. Let $\mu \in X(T)$ and $p > p(\mu)$. Then $H^p(Z, \psi^* \mathcal{L}(E(\mu))) = 0$.

As $\mathcal{O}_Z = \psi^*(\mathcal{O}_{G/B}) = \psi^* \mathcal{L}(E(0))$ we obtain by 4. Lemma (c) immediately the following

Corollary. $H^p(Z, \mathcal{O}_Z) = 0$ for all $p \geq 1$.

Proof of Theorem B. If E is a B -module, let $S(E^*) = \sum_q S_q(E^*)$ be the graded symmetrical algebra on the dual E^* of E . It may be considered as the ring of polynomial functions on E . The summands $S_q(E^*)$ are B -modules. Using [9] (9.4) one verifies that $\psi_* (\mathcal{O}_Z) = S((\mathcal{L}(\mathfrak{p}))^*) = \sum_q \mathcal{L}(S_q(\mathfrak{p}^*))$ and hence

$$\psi_* \psi^* \mathcal{L}(E(\mu)) = \psi_* (\mathcal{O}_Z) \otimes_{G/B} \mathcal{L}(E(\mu)) = \sum_q \mathcal{L}(S_q(\mathfrak{p}^*) \otimes_k E(\mu)).$$

Now it follows from [10] (1.3.3) and [8] Chapter II (3.10) that

$$H^p(Z, \psi^* \mathcal{L}(E(\mu))) = \sum_q H^p(S_q(\mathfrak{p}^*) \otimes_k E(\mu)).$$

Consider the $S(\mathfrak{g}^*)$ -module $M = S(\mathfrak{g}^*) \otimes_k E(\mu)$. Let J be the kernel of the canonical surjection $\mathfrak{g}^* \rightarrow \mathfrak{p}^*$. As x be a basis of J . As x is an M -regular sequence we have the long exact sequence

$$\dots K_2 \rightarrow K_1 \rightarrow K_0 \rightarrow S(\mathfrak{p}^*) \otimes_k E(\mu) \rightarrow 0$$

where $K_* = K(x; M)$ is the exterior complex, cf. [16] IV A2. Intrinsically the complex K_* may be defined by

$$K_n = M \otimes_k \Lambda^n J, \quad d_n: K_{n+1} \rightarrow K_n, \\ d_n(m \otimes (a_0 \wedge \dots \wedge a_n)) = \sum_{i=0}^n (-1)^i a_i m \otimes (a_0 \wedge \dots \wedge \hat{a}_i \wedge \dots \wedge a_n).$$

It is canonically graded by $K_* = \sum_q K^q$ where

$$K_n^q = S_{q-n}(\mathfrak{g}^*) \otimes_k E(\mu) \otimes \Lambda^n J.$$

Thus we have the long exact sequences of B -modules

$$\dots \rightarrow K_2^q \rightarrow K_1^q \rightarrow K_0^q \rightarrow S_q(\mathfrak{p}^*) \otimes_k E(\mu) \rightarrow 0.$$

As H^* is an exact delta-functor it suffices now to prove that $H^{p+n}(K_n^q) = 0$ for $q, n \geq 0$.

As $S_{q-n}(\mathfrak{g}^*)$ is a G -module we have the following cartesian square.

$$\begin{array}{ccc} G \times^B S_{q-n}(\mathfrak{g}^*) & \longrightarrow & S_{q-n}(\mathfrak{g}^*) \\ \downarrow & & \downarrow \\ G/B & \longrightarrow & G/G = pt. \end{array}$$

Considering $S_{q-n}(\mathfrak{g}^*)$ as a locally free sheaf on G/G we have therefore $\mathcal{L}(S_{q-n}(\mathfrak{g}^*)) = f^*(S_{q-n}(\mathfrak{g}^*))$ where $f: G/B \rightarrow G/G$ is the canonical morphism. By [10] (0_{III} 12.2.3) this implies

$$H^{p+n}(K_n^q) = S_{q-n}(\mathfrak{g}^*) \otimes H^{p+n}(E(\mu) \otimes \Lambda^n J).$$

So it suffices to prove that $H^{p+n}(E(\mu) \otimes \Lambda^n J) = 0$ for $n \geq 0$.

The B -module $E(\mu) \otimes \Lambda^n J$ has a filtration of B -modules F_i , $0 \leq i \leq r$, such that $F_0 = 0$ and $F_i/F_{i-1} \cong E(\chi_i)$ for some enumeration (χ_i) of the weights of $F_r = E(\mu) \otimes \Lambda^n J$. By convention the weights of \mathfrak{u} are the negative roots. Now dual modules have opposite weights and \mathfrak{p} contains \mathfrak{u} , so all positive roots are weights of \mathfrak{p}^* . Therefore the non-zero weights of J are negative roots with multiplicity one. Thus for every i there is a subset V_i of R_+ such that $\chi_i = \mu - |V_i|$ and $\#(V_i) \leq n$. If $w \in W$ is such that $w(\chi_i + \varrho)$ is dominant and regular, then we have

$$n(w) \leq p(\mu) + \#(V_i) < p + n.$$

By Bott's theorem as quoted in Section 3 this implies $H^{p+n}(E(\chi_i)) = 0$ for all i . It follows that $H^{p+n}(E(\mu) \otimes \Lambda^n J) = 0$.

References

1. Brieskorn, E.: Singular elements of semi-simple algebraic groups. Actes du Congrès intern. des mathématiciens 1970, t. II, 279—284
2. Burns, D.: On rational singularities in dimensions > 2 . Math. Ann. **211**, 237—244 (1974)
3. Séminaire H. Cartan, vol. 3, 1956—1957, New York: Benjamin 1967
4. Cartier, P.: Remarks on "Lie algebra cohomology and the generalized Borel-Weil theorem" by B. Kostant. Ann. Math. **74**, 388—390 (1961)
5. Chevalley, C.: Classification des groupes de Lie algébriques. Notes polycopiées, Inst. H. Poincaré, Paris 1956—1958
6. Demazure, M.: Une démonstration algébrique d'un théorème de Bott. Inventiones Math. **5**, 349—356 (1968)
7. Demazure, M.: Désingularisation des variétés de Schubert généralisées. Ann. Sc. Ec. Norm. Sup. 4^e série, t. 7, 53—88 (1974)
8. Godement, R.: Théorie des faisceaux. Paris: Hermann 1958
9. Grothendieck, A., Dieudonné, J. A.: Eléments de géométrie algébrique I. Grundlehren, 166. Berlin, Heidelberg, New York: Springer 1971
10. Grothendieck, A., Dieudonné, J. A.: Eléments de géométrie algébrique III. Publ. Math. de l'I.H.E.S. **11** (Paris 1961)
11. Hesselink, W. H.: Singularities in the nilpotent scheme of a classical group, to appear in Trans. A.M.S.
12. Kempf, G. et al.: Toroidal Embeddings I. Lecture Notes in Math. 339. Berlin, Heidelberg, New York: Springer 1973
13. Kempf, G. R.: Images of homogeneous vector bundles and varieties of complexes. Bull. A.M.S. **81**, 900—901 (1975)
14. Kostant, B.: Lie group representations on polynomial rings. Amer. J. Math. **85**, 327—404 (1963)
15. Macdonald, I. G.: The Poincaré series of a Coxeter group. Math. Ann. **199**, 161—174 (1972)
16. Serre, J.-P.: Algèbre locale—multiplicités. Lecture Notes in Math. 11. Berlin, Heidelberg, New York: Springer 1965
17. Springer, T. A.: The unipotent variety of a semisimple algebraic group. Proc. Bombay Coll. on Alg. Geometry 1968
18. Steinberg, R.: Conjugacy classes in algebraic groups. Lecture Notes in Math. 366. Berlin, Heidelberg, New York: Springer 1974

Received December 22, 1975